

Criteria-Based Consultation Prescribing Program

CRITERIA FOR DRUG COVERAGE

Testosterone 1% gel (Testim)

Initiation (new start) criteria: Non-formulary testosterone 1% gel (Testim) will be covered on the prescription drug benefit when the following criteria are met:

- All 5 of the following are met:
 1. Two low morning testosterone levels (below the normal range of the lab test used). The testosterone levels should be checked before testosterone treatment is started or 3 months after testosterone treatment has been stopped.
 - If body mass index (BMI) is greater or equal to 30, both total and free testosterone must be below the normal range. If the “free” testosterone is normal and the total testosterone is low, the diagnosis of hypogonadism cannot be made.
 2. Diagnosis of primary or secondary hypogonadism (*the gonads [testes in males] produce little or no sex hormones*); with documented luteinizing hormone (LH), follicle-stimulating hormone (FSH), and prolactin levels.
 3. If patient is over years of age 50: digital rectal exam (DRE) and prostate-specific antigen (PSA) test done in the last 12 months
 4. Hematocrit less than 50% (test that *measures the percent of red blood cells in the blood*) in the last 12 months.
 5. Adequate trial (at least 3 months) or intolerance/hypersensitivity with testosterone topical gel 1.62%.

-OR-

- Patient has a diagnosis of female to male gender dysphoria (*when a person’s gender identity does not match their biological sex*) AND intolerance/hypersensitivity with testosterone topical gel 1.62%.

-OR-

- Prescribed by a pediatric endocrinologist to cause puberty, or prior to genital surgery, or long-term puberty hormonal therapy AND intolerance/hypersensitivity with testosterone topical gel 1.62%.

-OR-

kp.org

Revised: 09/09/21
Effective: 12/02/21

All plans offered and underwritten by
Kaiser Foundation Health Plan of the Northwest

Criteria-Based Consultation Prescribing Program

CRITERIA FOR DRUG COVERAGE

Testosterone 1% gel (Testim)

- Prescribed by Obstetrics/Gynecology or Urology for post-menopausal hyposexual drive disorder/dysDD).
- Note: Testim is covered under the prescription drug benefit for HSDD only for members with coverage for medications used to treat sexual dysfunction. Others pay member cash price.

Criteria for members already taking the medication who have not been reviewed previously (e.g., new members): Non-formulary testosterone 1% gel (Testim) will be covered on the prescription drug benefit when the following criteria are met:

- All 5 of the following are met:
 1. Two low morning testosterone levels (below the normal range of the lab test used). The testosterone levels should be checked before testosterone treatment is started or 3 months after testosterone treatment has been stopped.
 - If body mass index (BMI) is greater or equal to 30, both total and free testosterone must be below the normal range. If the “free” testosterone is normal and the total testosterone is low, the diagnosis of hypogonadism cannot be made.
 2. Diagnosis of primary or secondary hypogonadism (*the gonads [testes in males] produce little or no sex hormones*); with documented luteinizing hormone (LH), follicle-stimulating hormone (FSH), and prolactin levels.
 3. If patient is over years of age 50: digital rectal exam (DRE) and prostate-specific antigen (PSA) test done in the last 12 months.
 4. Hematocrit less than 50% (test that *measures the percent of red blood cells in the blood*) in the last 12 months.
 5. Adequate trial (at least 3 months) or intolerance/hypersensitivity with testosterone topical gel 1.62%.

-OR-

- Patient has a diagnosis of female to male gender dysphoria (*when a person’s gender identity does not match their biological sex*) AND intolerance/hypersensitivity with testosterone topical gel 1.62%.

kp.org

Revised: 09/09/21
Effective: 12/02/21

All plans offered and underwritten by
Kaiser Foundation Health Plan of the Northwest

Criteria-Based Consultation Prescribing Program CRITERIA FOR DRUG COVERAGE

Testosterone 1% gel (Testim)

-OR-

- Prescribed by a pediatric endocrinologist to cause puberty, or prior to genital surgery, or long-term puberty hormonal therapy AND intolerance/hypersensitivity with testosterone topical gel 1.62%.

-OR-

- Prescribed by or in consultation with Obstetrics/Gynecology or Urology for post-menopausal hyposexual drive disorder/dysfunction (HSDD).
- Note: Testim is covered under the prescription drug benefit for HSDD only for members with coverage for medications used to treat sexual dysfunction. Others pay member cash price.

kp.org

Revised: 09/09/21
Effective: 12/02/21

All plans offered and underwritten by
Kaiser Foundation Health Plan of the Northwest