



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

Generic	Brand	HICL	GCN	Medi-Span	Exception/Other
TESTOSTERONE	ANDRODERM, ANDROGEL, AXIRON, FORTESTA, NATESTO, STRIANT, TESTIM, VOGELXO	01403		GPI-10 (2310003000)	BRAND ≠ TESTOPEL ROUTE ≠ MISCELL., IMPLANT
TESTOSTERONE CYPIONATE	DEPO- TESTOSTERONE	01400		GPI-10 (2310003010)	NDC ≠ 76420065001 FDB: ROUTE ≠ MISCELL. MEDISPAN: ROUTE ≠ DOES NOT APPLY.
TESTOSTERONE ENANTHATE	TESTOSTERONE ENANTHATE, XYOSTED	01401		GPI-10 (2310003020)	FDB: ROUTE ≠ MISCELL.
METHYLTESTOSTERONE	TESTRED, ANDROID, METHITEST		10380 10411	GPI-10 (2310002000)	
TESTOSTERONE UNDECANOATE	JATENZO	07304		GPI-10 (2310003080)	BRAND ≠ AVEED

GUIDELINES FOR USE

INITIAL CRITERIA (NOTE: FOR RENEWAL CRITERIA SEE BELOW)

1. Is the request for a male patient with a diagnosis of primary or secondary hypogonadism (hypotestosteronism or low testosterone) who meets **ONE** of the following criteria?
 - The patient has a previously approved prior authorization for testosterone or has been receiving any form of testosterone replacement therapy as indicated per physician attestation or claims history **OR**
 - The patient has **AT LEAST ONE** of the following laboratory values confirming low testosterone levels:
 - At least two morning total serum testosterone levels of less than 300 ng/dL (10.4 nmol/L) taken on separate occasions while in a fasted state
 - Free serum testosterone level of less than 5 pg/mL (0.17 nmol/L)

If yes, continue to #2.

If no, continue to #7.

CONTINUED ON NEXT PAGE

Copyright © 2021 MedImpact Healthcare Systems, Inc. All rights reserved. This document is proprietary to MedImpact. MedImpact maintains the sole and exclusive ownership, right, title, and interest in and to this document.



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

INITIAL CRITERIA (CONTINUED)

2. Is the request for Xyosted **AND** have the following criteria been met?

- The patient is 18 years of age or older
- The requested medication is being used for testosterone replacement therapy

If yes, **approve the requested strength for 12 months by GPID or GPI-10 with a quantity limit of #4 syringes per 28 days.**

If no, continue to #3.

3. Is the request for Jatenzo **AND** has the following criterion been met?

- The patient is 18 years of age or older

If yes, **approve for 12 months by GPID or GPI-14 for all strengths with the following quantity limits:**

- **Jatenzo 158mg: #4 per day.**
- **Jatenzo 198mg: #4 per day.**
- **Jatenzo 237mg: #2 per day.**

If no, continue to #4

4. Is the request for AndroGel 1%, AndroGel 1.62%, Axiron, Testim, Vogelxo, Depo-Testosterone (testosterone cypionate), or intramuscular testosterone enanthate?

If yes, **approve the requested agent for 12 months by GPID or GPI-14 with the following quantity limits:**

- **AndroGel 1% (testosterone) (2.5 gram packet): #5 grams per day; (5 gram packet): #10 grams per day; (75 gram pump): #300 grams (4 pumps) per 30 days.**
- **AndroGel 1.62% (testosterone) (1.25 gram packet): #1.25 grams per day; (2.5 gram packet): #5 grams per day; (75 gram pump): #150 grams (2 pumps) per 30 days.**
- **Axiron (testosterone) (90 mL pump): #180 mL per 30 days.**
- **Testim (testosterone) (5 gram gel tube): #10 grams per day.**
- **Vogelxo (testosterone) (5 gram gel tube): #10 grams per day; (5 gram gel packet): #10 grams per day; (75 gram pump): #300 grams (4 pumps) per 30 days.**
- **Depo-Testosterone (testosterone cypionate) (100mg/mL, 200mg/mL [10mL vial]): up to #10mL per 28 days.**
- **Depo-Testosterone (testosterone cypionate) (200mg/mL [1mL vial]): up to #10mL per 30 days.**
- **Intramuscular testosterone enanthate (200mg/mL [5mL vial]): #5mL per 28 days.**

If no, continue to #5.

CONTINUED ON NEXT PAGE



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

INITIAL CRITERIA (CONTINUED)

5. Is the request for Androderm patches, Fortesta, Natesto, or Striant, **AND** has the following criterion been met?

- The patient had a trial of or contraindication to a generic lower cost agent (e.g., AndroGel 1%, AndroGel 1.62%, Axiron, Testim, Vogelxo, Depo-Testosterone, intramuscular testosterone enanthate)

If yes, **approve the requested agent for 12 months by GPID or GPI-14 with the following quantity limits:**

- **Androderm (testosterone) (2mg, 4mg patches): #30 patches per 30 days.**
- **Fortesta (testosterone) (60 gram pump): #120 grams (2 pumps) per 30 days.**
- **Natesto (testosterone) (7.32 gram bottle): #21.96 grams (3 bottles) per 30 days.**
- **Striant (testosterone): #60 buccal systems per 30 days.**

If no, continue to #6.

6. Is the request for Android, Methitest, or Testred, **AND** has the following criterion been met?

- The patient had a trial of or contraindication to **TWO** lower cost agents (e.g., AndroGel 1%, Axiron, Testim, Vogelxo, Depo-Testosterone, intramuscular testosterone enanthate, Androderm, AndroGel 1.62%, Fortesta, Natesto, Striant, Jatenzo)

If yes, **approve the requested agent for 12 months by GPID or GPI-14 with the following quantity limits:**

- **Android/Testred (methyltestosterone) (10mg capsule): #5 capsules per day.**
- **Methitest (methyltestosterone) (10mg tablet): #5 tablets per day.**

If no, do not approve.

DENIAL TEXT: See the initial denial text at the end of the guideline.

CONTINUED ON NEXT PAGE



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

INITIAL CRITERIA (CONTINUED)

7. Is the request for a male patient with a diagnosis of delayed puberty not secondary to a pathological disorder and meets **ONE** of the following criteria?

- The request is for intramuscular testosterone enanthate
- The request is for methyltestosterone (Testred, Android, or Methitest) **AND** the patient had a previous trial of or contraindication to intramuscular testosterone enanthate

If yes, **approve the requested agent for lifetime by GPID or GPI-14 with the following quantity limits:**

- **Intramuscular testosterone enanthate (200mg/mL, 5mL vial): #5mL per 28 days.**
- **Testred/Android (methyltestosterone) (10mg capsule): #5 capsules per day.**
- **Methitest (methyltestosterone) (10mg tablet): #5 tablets per day.**

If no, continue to #8.

8. Is the requested agent for gender dysphoria as supported by the compendia (e.g., DrugDex strength of recommendation Class I, IIa, or IIb) **AND** has the following criterion been met?

- The patient is 16 years of age or older

If yes, **approve the requested agent for 12 months by GPID or GPI-14 and override quantity limits.**

If no, continue to #9.

9. Is the request for a female patient with a diagnosis of metastatic breast cancer and meets **ONE** of the following criteria?

- The request is for intramuscular testosterone enanthate
- The request is for methyltestosterone (Testred, Android, or Methitest) **AND** the patient had a previous trial of or contraindication to intramuscular testosterone enanthate

If yes, **approve the requested agent for lifetime by GPID or GPI-14 with the following quantity limits:**

- **Intramuscular testosterone enanthate (200mg/mL, 5mL vial): #5mL per 28 days.**
- **Testred/Android (methyltestosterone) (10mg capsule): #20 capsules per day.**
- **Methitest (methyltestosterone) (10mg tablet): #20 tablets per day.**

If no, do not approve.

DENIAL TEXT: See the initial denial text at the end of the guideline.

CONTINUED ON NEXT PAGE



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

INITIAL CRITERIA (CONTINUED)

INITIAL DENIAL TEXT: *Some terms are already pre-defined in parenthesis. Please use these definitions if the particular text you need to use does not already have definition(s) in it.

Our guideline named **TESTOSTERONE** requires the following rule(s) be met for approval:

- A. You have ONE of the following diagnoses:
 - 1. Primary or secondary male hypogonadism (hypotestosteronism or low testosterone)
 - 2. Delayed puberty in males not due to a pathological disorder (not due to disease)
 - 3. Gender dysphoria (you identify yourself as a member of the opposite sex)
 - 4. Female with metastatic breast cancer (cancer spreading to other areas of body)
- B. **If you are a female with metastatic breast cancer or you are a male with delayed puberty not secondary to a pathological (extreme) disorder**, only intramuscular (injected into muscle) testosterone enanthate or methyltestosterone (Testred, Android, or Methitest) may be approved
- C. **If you have gender dysphoria, approval also requires:**
 - 1. Only agents supported by the compendia (accepted medical references such as DrugDex strength of recommendation Class I, IIa, or IIb) for treatment of gender dysphoria may be approved
 - 2. You are 16 years of age or older
- D. **If you are a male with primary or secondary hypogonadism, approval requires ONE of the following:**
 - 1. You have a previously approved prior authorization for testosterone, or you have been receiving any form of testosterone replacement therapy as indicated per physician attestation or claims history OR
 - 2. You have ONE of the following lab values showing you have low testosterone levels:
 - i. At least two morning total serum (blood) testosterone levels of less than 300 ng/dL (10.4 nmol/L) taken on separate occasions while in a fasted state (you have not eaten)
 - ii. Free serum testosterone level of less than 5 pg/mL (0.17 nmol/L)
- E. **If the request is for Xyosted, approval also requires:**
 - 1. You are 18 years of age or older
 - 2. The requested medication is being used for testosterone replacement therapy
- F. **If the request is for Jatenzo, approval also requires:**
 - 1. You are 18 years of age or older
- G. **If the request is for Androderm, Fortesta, Natesto or Striant, approval also requires:**
 - 1. You had a trial of a generic lower cost agent (e.g. AndroGel 1%, AndroGel 1.62%, Axiron, Testim, Vogelxo, Depo-Testosterone, intramuscular testosterone enanthate), unless there is a medical reason why you cannot (contraindication)

(Initial denial text continued on next page)

CONTINUED ON NEXT PAGE



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

INITIAL CRITERIA (CONTINUED)

H. If the request is for Android, Methitest, or Testred, approval also requires:

1. You had a trial of **TWO** lower cost agents (e.g. AndroGel 1%, Axiron, Testim, Vogelxo, Depo-Testosterone, intramuscular (injected into the muscle) testosterone enanthate, Androderm, AndroGel 1.62%, Fortesta, Natesto, Striant, Jatenzo), unless there is a medical reason why you cannot (contraindication)

I. If you are a male patient requesting methyltestosterone (Testred, Android or Methitest) for delayed puberty not secondary to a pathological disorder, approval also requires:

1. You had a previous trial of intramuscular (injected into the muscle) testosterone enanthate, unless there is a medical reason why you cannot (contraindication). Please note that Intramuscular testosterone enanthate requires a prior authorization

J. If you are a female patient requesting methyltestosterone (Testred, Android or Methitest) for metastatic breast cancer, approval also requires:

1. You had a previous trial of intramuscular (injected into the muscle) testosterone enanthate, unless there is a medical reason why you cannot (contraindication). Please note that intramuscular testosterone enanthate requires a prior authorization.

Your doctor told us [INSERT PT SPECIFIC INFO PROVIDED]. We do not have information showing you [INSERT UNMET CRITERIA]. This is why your request is denied. Please work with your doctor to use a different medication or get us more information if it will allow us to approve this request.

CONTINUED ON NEXT PAGE



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

INITIAL CRITERIA (CONTINUED)

RENEWAL CRITERIA

1. Is the request for a male patient with a diagnosis of primary or secondary hypogonadism (hypotestosteronism or low testosterone) who meets **ALL** of the following criteria?
 - The patient has improved symptoms compared to baseline and tolerance to treatment
 - Documentation of normalized serum testosterone levels and hematocrit concentrations compared to baseline

If yes, **approve requested agent for 12 months by GPID or GPI-14 with the following quantity limits:**

- **Xyosted (testosterone enanthate) (50mg/0.5mL, 75mg/0.5mL, 100mg/0.5mL subcutaneous auto-injectors): #4 syringes per 28 days.**
- **Jatenzo (testosterone undecanoate): (158 mg and 198 mg): #4 per day; (237 mg): #2 per day.**
- **AndroGel 1% (testosterone) (2.5 gram packet): #5 grams per day; (5 gram packet): #10 grams per day; (75 gram pump): #300 grams (4 pumps) per 30 days.**
- **Axiron (testosterone) (90 mL pump): #180 mL per 30 days.**
- **Testim (testosterone) (5 gram gel tube): #10 grams per day.**
- **Vogelxo (testosterone) (5 gram gel tube): #10 grams per day; (5 gram gel packet): #10 grams per day; (75 gram pump): #300 grams (4 pumps) per 30 days.**
- **Depo-Testosterone (testosterone cypionate): (100mg/mL, 200mg/mL [10mL vial]): up to #10mL per 28 days.**
- **Depo-Testosterone (testosterone cypionate): (200mg/mL [1mL vial]): up to #10mL per 30 days.**
- **Intramuscular testosterone enanthate (200mg/mL, 5mL vial): #5mL per 28 days.**
- **Androderm (testosterone) (2mg, 4mg patches): #30 patches per 30 days.**
- **AndroGel 1.62% (testosterone) (1.25 gram packet): #1.25 grams per day; (2.5 gram packet): #5 grams per day; (75 gram pump): #150 grams (2 pumps) per 30 days.**
- **Fortesta (testosterone) (60 gram pump): #120 grams (2 pumps) per 30 days.**
- **Natesto (testosterone) (7.32 gram bottle): #21.96 grams (3 bottles) per 30 days.**
- **Striant (testosterone) #60 buccal systems per 30 days.**
- **Android (methyltestosterone) (10mg capsule): #5 capsules per day.**
- **Methitest (methyltestosterone) (10mg tablet): #5 tablets per day.**
- **Testred (methyltestosterone) (10mg capsule): #5 capsules per day.**

If no, continue to #2.

CONTINUED ON NEXT PAGE



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

RENEWAL CRITERIA (CONTINUED)

2. Is the request for a male patient with a diagnosis of delayed puberty not secondary to a pathological disorder?

If yes, **approve the requested agent for lifetime by GPID or GPI-14 with the following quantity limits:**

- **Intramuscular testosterone enanthate (200mg/mL, 5mL vial): #5mL per 28 days.**
- **Testred/Android (methyltestosterone) (10mg capsule): #5 capsules per day.**
- **Methitest (methyltestosterone) (10mg tablet): #5 tablets per day.**

If no, continue to #3.

3. Is the request for a female patient with a diagnosis of metastatic breast cancer?

If yes, **approve the requested agent for lifetime by GPID or GPI-14 with the following quantity limits:**

- **Intramuscular testosterone enanthate (200mg/mL, 5mL vial): #5mL per 28 days.**
- **Testred/Android (methyltestosterone) (10mg capsule): #20 capsules per day.**
- **Methitest (methyltestosterone) (10mg tablet): #20 tablets per day.**

If no, continue to #4.

4. Is the requested agent for gender dysphoria as supported by the compendia (e.g. DrugDex strength of recommendation Class I, IIa, or IIb)?

If yes, **approve the requested agent for 12 months by GPID or GPI-14 and override quantity limits.**

If no, do not approve.

DENIAL TEXT: See the renewal denial text at the end of the guideline.

CONTINUED ON NEXT PAGE



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

INITIAL CRITERIA (CONTINUED)

RENEWAL DENIAL TEXT: *Some terms are already pre-defined in parenthesis. Please use these definitions if the particular text you need to use does not already have definition(s) in it.

Our guideline named **TESTOSTERONE** requires the following rule(s) be met for renewal:

- A. You have ONE of the following diagnoses:
 - 1. Primary or secondary male hypogonadism (hypotestosteronism or low testosterone)
 - 2. Delayed puberty in males not due to a pathological (extreme) disorder (not due to disease)
 - 3. Female with metastatic breast cancer (cancer spreading to other areas of body)
 - 4. Gender dysphoria (you identify yourself as a member of the opposite sex)
- B. **If you have gender dysphoria, renewal also requires:**
 - 1. Only agents supported by the compendia (accepted medical references such as DrugDex strength of recommendation Class I, IIa, or IIb) for treatment of gender dysphoria may be approved
- C. **If you are a male patient with primary or secondary hypogonadism, renewal also requires:**
 - 1. You have shown improvement in your symptoms compared to baseline and tolerance to treatment
 - 2. Documentation of normalized serum testosterone levels and hematocrit concentrations (type of blood test) compared to baseline
- D. **If you are a male patient with delayed puberty not secondary to a pathological disorder, only the following medications will be approved:**
 - 1. Intramuscular testosterone enanthate, Testred, Android, Methitest
- E. **If you are a female patient with metastatic breast cancer, only the following medications will be approved:**
 - 1. Intramuscular testosterone enanthate, Testred, Android, Methitest

Your doctor told us **[INSERT PT SPECIFIC INFO PROVIDED]**. We do not have information showing you **[INSERT UNMET CRITERIA]**. This is why your request is denied. Please work with your doctor to use a different medication or get us more information if it will allow us to approve this request.

CONTINUED ON NEXT PAGE



STANDARD COMMERCIAL DRUG FORMULARY
PRIOR AUTHORIZATION GUIDELINES

TESTOSTERONE

RATIONALE

For further information, please refer to the Prescribing Information and/or Drug Monograph for the related testosterone formulation.

REFERENCES

- Androderm [Prescribing Information]. Parsippany, NJ: Allergan; May 2020.
- Android [Prescribing Information]. Bridgewater, NJ: Valeant Pharmaceuticals.; April 2015.
- Androgel 1% [Prescribing Information]. North Chicago, IL: AbbVie Inc.; April 2020.
- Androgel 1.62% [Prescribing Information]. North Chicago, IL: Abbvie Inc.; April 2020.
- Axiron [Prescribing Information]. Indianapolis, IN: Lilly USA, LLC.; July 2017.
- Depo-Testosterone [Prescribing Information]. New York, NY: Pharmacia & Upjohn Company.; August 2020.
- Fortesta [Prescribing Information]. Malvern, PA: Endo Pharmaceuticals.; June 2020.
- Methitest [Prescribing Information]. Hayward, CA: Impax Generics.; May 2019.
- Natesto [Prescribing Information]. Englewood, CO: Aytu BioScience Inc.; December 2019.
- Striant [Prescribing Information]. Malvern, PA: Actient Pharmaceuticals LLC.; October 2016.
- Testim [Prescribing Information]. San Antonio, TX: DPT Laboratories, Ltd.; May 2019.
- Testred [Prescribing Information]. Bridgewater, NJ: Valeant Pharmaceuticals.; April 2015.
- Vogelxo [Prescribing Information]. Maple Grove, MN: Upsher-Smith Lab., Inc.; July 2020.
- Xyosted [Prescribing Information]. Ewing, NJ: Antares Pharma Inc.; November 2019.
- Jatenzo [Prescribing Information]. Northbrook, IL: Clarus Therapeutics, Inc.; September 2019.

Library	Commercial	NSA
Yes	Yes	No

Part D Effective: N/A

Commercial Effective: 01/01/21

Created: 02/01

Client Approval: 11/20

P&T Approval: 01/20